

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) Publication number:

0 504 202 B1

(12)

EUROPEAN PATENT SPECIFICATION

(49) Date of publication of patent specification: 03.05.95 (51) Int. Cl.⁶: **A61K 39/095, A61K 39/108, A61K 39/385**

(21) Application number: **91900142.0**

(22) Date of filing: **13.12.90**

(86) International application number:
PCT/CA90/00437

(87) International publication number:
WO 91/08772 (27.06.91 91/14)

(54) **IMPROVED MENINGOCOCCAL POLYSACCHARIDE CONJUGATE VACCINE.**

(30) Priority: **14.12.89 US 448195**

(43) Date of publication of application:
23.09.92 Bulletin 92/39

(45) Publication of the grant of the patent:
03.05.95 Bulletin 95/18

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(56) References cited:
EP-A- 98 581
US-A- 4 057 685
US-A- 4 356 170
US-A- 4 727 136

(73) Proprietor: **NATIONAL RESEARCH COUNCIL OF CANADA**
Montreal Road
Ottawa
Ontario K1A 0R6 (CA)

(72) Inventor: **JENNINGS, Harold, J.**
2049 Woodglen Crescent
Gloucester, ON, K1J 6G6 (CA)
Inventor: **MICHON, Francis**
128 Keefer Street
Ottawa, ON, K1M 1T5 (CA)

(74) Representative: **Laufhütte, Dieter, Dr.-Ing. et al**
Lorenz-Seldler-Gossel
Widenmayerstrasse 23
D-80538 München (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 504 202 B1

Description

The present invention is directed to chemically-modified group B polysaccharides of *Neisseria meningitidis*. The invention also provides vaccines in which the respective modified polysaccharides are conjugated to a protein carrier.

BACKGROUND OF THE INVENTION

Meningitis caused by group B *N. meningitidis* and *E. coli* K1 remain major world health problems. Group B meningitis occurs in both endemic and epidemic situations and accounts for approximately half of all recorded cases of meningococcal meningitis, while K1-positive *E. coli* are the leading cause of meningitis in neonates. Currently there is no vaccine commercially available against disease caused by group B meningococci and *E. coli* K1. This is in large part due to the fact that the group B meningococcal polysaccharide (GBMP) is only poorly immunogenic in humans. There are some recently reported candidate vaccines based on complexes of the GBMP with outer membrane proteins, but, as yet, there is no clear evidence of their efficacy in humans.

Recently, a new concept of a vaccine based on a synthetic chemically modified (N-propionylated) group B polysaccharide-protein (N-Pr-GBMP-protein) conjugate has been developed. The vaccine induces in mice high titers of IgG antibodies which are not only protective, but also cross-react with unmodified GBMP (i.e. N-acetyl-GBMP). This concept is described and claimed in U.S. patent No. 4,727,136, issued February 23, 1988 to Harold J. Jennings et al.

It has been inferred that a vaccine which raises cross-reactive antibodies such as that described in U.S. patent 4,727,136 could only be successful at the expense of breaking immune tolerance. This hypothesis is legitimized by the identification of a common epitope consisting of a chain of α -(2-8)-linked sialic acid residues (with a minimum requirement of ten residues) in both the native N-Ac-GBMP and in human and animal tissue (Jennings, *Contrib. Microbiol. Immunol.* Basel, Karger, 1989, Vol. 10, 151-165). These polysialosyl chains function as developmental antigens and have for the most part been associated with the fetal state in embryonic neural cell adhesion (Finne et al, *Biochem. Biophys. Res. Commun.*, 1983, 112, 482). During post-natal maturation, this antigen is down-regulated (Friedlander et al, *J. Cell Biol.* 1985, 101, 412) but is expressed in mature humans during the regeneration of diseased muscles (Cashman et al, *Ann. Neuron.*, 1987, 21, 481) in tumor cells (Roth et al, *Proc. Natl. Acad. Sci.*, 1988, 85, 299) and in natural killer (NK) and CD3⁺T cells (Husmann et al, *Eur. J. Immunol.*, 1989, 19, 1761). Although the consequences of breaking tolerance to these fetal antigens have not yet been established, it is generally conceded that, because of this cross-reaction, the N-Pr-GBMP-protein conjugate would be severely scrutinized by licensing agencies, resulting in considerable expense and delays because of the complex experimentation necessary to prove the safety of the vaccine before its approval for commercial marketing.

It is an object of the present invention to develop a vaccine having immunogenic properties which are enhanced as compared to those of the N-Pr-GBMP-protein. It is also an object of the invention to provide a vaccine which exhibits substantially reduced cross-reactivity with a human and animal neural cell adhesion antigen (fetal N-CAM).

Summary of the Invention

In one aspect of the present invention, there is provided a modified B polysaccharide of *Neisseria meningitidis* having sialic acid residue N-acetyl (C₂) groups replaced by a C₄ - C₈ acyl group, wherein the modified polysaccharide has an average molecular weight within the range of 10,000 to 50,000 Daltons.

In another aspect, there is provided an antigenic conjugate comprising the N-C₄-C₈ acyl polysaccharide conjugated to an immunologically suitable protein, wherein the modified B polysaccharide of *Neisseria meningitidis* has an average molecular weight within the range of 10,000 to 50,000 Daltons, having enhanced immunogenicity with substantially reduced inducement of cross-reactive antibodies.

In a further aspect, there is provided a vaccine comprising said N-C₄-C₈ acyl polysaccharide-protein conjugate in association with a suitable carrier or diluent. The vaccines of the invention may also comprise a therapeutically effective amount of an adjuvant suitable for human use, for example aluminum phosphate or aluminum hydroxide.

In yet another aspect, there is provided a modified *E. coli* K1 capsular polysaccharide wherein the sialic acid residue N-acetyl groups of the native polysaccharide are replaced by a C₄-C₈ acyl group wherein the modified polysaccharide has an average molecular weight within the range of 10,000 to 50,000 Daltons.

In a yet further aspect, there is provided a gamma globulin fraction capable of passive protection against meningitis caused by group B *N. meningitidis* and *E. coli* K1, said gamma globulin fraction being produced by immunizing a mammal with said vaccine and recovering gamma globulin from said mammal.

The fraction is produced by immunizing a mammal with a vaccine of the invention. The fraction is then administered to an individual to provide protection against or to treat on-going infection caused by the above organisms. From this, it will be appreciated that the immunogenic vaccine conjugates of the invention will provide for a source of therapeutic antiserum in light of their favorable immunogenicity with minimal inducement of antibodies cross-reactive with the human and animal neural cell adhesion antigen fetal N-CAM. The conjugates of the invention will also be useful for raising monoclonal antibodies and, possibly, antidiotype antibodies.

It has been found in our recent experiments that most of the bactericidal and protective antibodies induced by the N-Pr-GBMP-protein conjugate described in the above-referred to Jennings et al U.S. patent 4,727,136 are not associated with the fetal N-CAM cross-reactive antibodies. In fact, most of the protective activity is contained in an N-Pr-GBMP-specific antibody population which does not cross-react with fetal N-CAM. In light of this, it is believed that the N-Pr-GBMP mimics a unique bactericidal epitope on the surface of group B meningococci.

The present invention is based on the discovery that it is possible to synthesize chemically modified GBMP's which mimic the bactericidal epitope and which, in their conjugated form, not only exhibit enhanced immunogenicity but also avoid substantially the inducement of antibodies that cross-react with fetal N-CAM.

In arriving at the present invention, a number of different chemically modified GBMP's have been synthesized and conjugated individually to protein, followed by injection of the conjugates into mice and the effects compared to those produced by the N-Pr-GBMP protein conjugate. Surprisingly, it has been found that the N-C₄-C₈ acyl GBMP-protein conjugates, for example the n-butanoyl, iso-butanoyl, n-pentanoyl, iso-pentanoyl, neo-pentanoyl, hexanoyl, heptanoyl, and octanoyl, and especially the N-butanoyl (N-Bu) GBMP-protein conjugate, substantially mimic the bacterial epitope with substantially reduced inducement of cross-reactive antibodies.

DETAILED DESCRIPTION OF THE INVENTION

The group B meningococcal polysaccharide is isolated from *N. meningitidis* by methods which are known in the art. In one such method, group B meningococci (strain 981B) were grown at 37°C in a fermenter using 30 g. of dehydrated Todd Hewitt Broth (Difco Laboratories, Detroit, Michigan) per liter of distilled water. Prior to fermenter growth, the lyophilized strain was grown initially in a candle jar at 37°C on 5% (v/v) Sheep's Blood Agar (Difco Laboratories, Detroit, Michigan) plates. The bacteria were then transferred to 1.0 liter of Todd Hewitt Broth (as above) in an Erlenmeyer flask which was shaken at 37°C for 7 hours at 190 r.p.m. This inoculum was then transferred to the fermenter. After fermenter growth (16 hours) the bacteria were killed by the addition of formalin to a final concentration of 0.75%. The bacteria were removed by continuous centrifugation and the group B meningococcal polysaccharide was isolated from the supernatant and purified essentially as described by Bundle et al, *J. Biol. Chem.*, 249, 4797-4801 (1974) except that the protein was extracted by stirring a solution of the crude polysaccharide with cold (4°C) 90% phenol instead of hot (50-60°C). This latter process ensures that a high molecular weight form of the GBMP is produced.

E. coli (O18:K1:H7) (NRCC 4283) were grown at 37°C in a fermenter in distilled water containing dehydrated Brain Heart Infusion (BHI; 37 g/litre) (Difco Laboratories, Detroit, Michigan). Prior to fermenter growth, the lyophilized strain was grown on 50 ml of BHI solution (same as above) in an Erlenmeyer flask which was shaken at 37°C for 7 hours at 200 r.p.m. This growth was then transferred to 1.5 litres of BHI (as above) and grown under the same conditions as described above for 7 hours. The inoculum was then transferred to the fermenter.

The procedures employed in the isolation and purification of the capsular polysaccharide of *E. coli* K1 were identical to those described above for the isolation of the group B meningococcal polysaccharide.

It will be appreciated that the isolation and purification procedures described above are not the only ones which may be utilized, and that other published procedures are available, for example those described by Watson et al, *J. Immunol.*, 81, 331 (1958) and in the above-mentioned U.S. patent 4,727,136.

The native polysaccharide is N-deacetylated to provide a reactive amine group in the sialic acid residue parts of the molecule. The N-deacetylation can be carried out by any known method, for example in a basic aqueous medium at elevated temperatures, for example about 90° to 110°C., and at a pH of about 13 to 14. The basic aqueous medium is suitably an aqueous alkali metal hydroxide solution, for example sodium

hydroxide of about 2M concentration. Alternatively, hydrazine in aqueous solution may be used. The degree of N-deacetylation may vary from about 30% to 100%, depending on the conditions. It is preferred to achieve about 90 to 100% N-deacetylation. N-deacetylated product can be recovered for example by cooling, neutralizing, purification if desired, and lyophilization.

5 Prior to the N-deacetylation procedure, the native polysaccharide has an average molecular weight in the region of about 500,000 to 800,000 Daltons. As a result of N-deacetylation, fragments of the polysaccharide are produced having an average molecular weight ranging from about 10,000 to 50,000 Daltons.

The N-deacetylated polysaccharide fragments are then N-acylated to produce the corresponding N-acylated product. The N-acylation may be carried out by dissolving the N-deacetylated polysaccharide in an aqueous medium having a pH of about 7.5 to 9.0, followed by adding the appropriate acyl anhydride, optionally with an alcohol to increase solubility, and cooling to below 10°C until the reaction is complete. The reaction medium can be purified, if desired, for example by dialysis, and the N-acylated product then recovered, typically by lyophilization. The reaction is substantially complete within about 10 to 20 hours. 15 The degree of N-acylation, as measured by analytical techniques, typically ¹H nmr, is at least 90%, and likely close to 100%. The N-acylation reaction does not result in any significant molecular weight reduction of the fragments.

It is preferred, according to the present invention, to select, for conjugation purposes, the N-acylated material having an average molecular weight corresponding to about 30 to 200 sialic acid residues. This is generally achieved by way of gel filtration of the N-acylated GBMP using an Ultragel (trademark) AcA 44 (Bead diameter 60-140 µm) column, using PBS as eluant. Alternatively, a suitable sizing membrane may be employed.

N-acylated material of average molecular weight of 10,000 to 50,000 Daltons, for example 10,000 to 15,000 Daltons, is employed for the invention. This is obtained by collecting the fractions of the eluate of the column containing N-acylated GBMP material having that average molecular weight range. N-acylated material of higher average molecular weight, for example in the region of 30,000 to 40,000 Daltons, has also proved to be useful according to the invention. 25

The vaccines of the invention are produced by conjugating the N-acylated polysaccharide with an immunologically suitable carrier protein. Preferably, the carrier protein itself is an immunogen. Examples of suitable carrier proteins are tetanus toxoid, diphtheria toxoid, cross-reacting materials (CRMs), preferably CRM₁₉₇, obtained from Sclavo Ltd., Siena, Italy, and bacterial protein carriers, for example meningococcal outer membrane proteins. 30

Any mode of conjugation may be employed to conjugate the modified polysaccharide fragments with the carrier protein. A preferred method is that described in U.S. patent 4,356,170, i.e. by introducing terminal aldehyde groups (via oxidation of cis-vicinal hydroxyl groups) into the N-acylated polysaccharide and coupling the aldehyde groups to the protein amino groups by reductive amination. The polysaccharide and the protein are thereby linked through a -CH₂-NH-protein linkage. 35

It is to be understood, however, that the conjugate vaccines of the invention are not limited to those produced via reductive amination. Thus, the vaccines may also be produced by conjugating the N-acylated polysaccharide with the carrier protein using an adipic dihydrazide spacer, as described by Schneerson, R., et al, Preparation, Characterization and Immunogenicity of Haemophilus influenzae type b Polysaccharide-Protein Conjugates, J. Exp. Med., 1952, 361-476 (1980), and in U.S. patent 4,644,059 to Lance K. Gordon. Alternatively, there may be used the binary spacer technology developed by Merck, as described by Marburg, S., et al, "Biomolecular Chemistry of Macromolecules: Synthesis of Bacterial Polysaccharide Conjugates with Neisseria meningitidis Membrane Protein", J. Am. Chem. Soc., 108, 5282-5287 (1986) or, 45 possibly, the reducing ends methodology, as referred to by Anderson in U.S. patent 4,673,574.

The resulting N-acylated polysaccharide-protein conjugates do not possess significant cross-linking and are soluble in aqueous solutions. This makes the conjugates of the invention good candidates for vaccine use.

50 The resulting N-acylate-dpolysaccharide-protein conjugates of the invention have been tested in *in vitro* tests in mice, and have generally been shown to possess improved immunogenic properties as compared with the N-propionylated-polysaccharide. In addition, substantially reduced formation of cross-reactive antibodies is observed. In light of this, it is believed that the vaccines of the invention will be useful against meningitis caused by group B *N. meningitidis* or by *E. coli* K1 organisms. Of particular interest are vaccines for protecting human infants who are most susceptible to bacterial meningitis. 55

The vaccines of the invention are typically formed by dispersing the conjugate in any suitable pharmaceutically acceptable carrier, such as physiological saline or other injectable liquids. The vaccine is administered parenterally, for example subcutaneously, intraperitoneally or intramuscularly. Additives cus-

tomary in vaccines may also be present, for example stabilizers such as lactose or sorbitol and adjuvants such as aluminum phosphate, hydroxide, or sulphate.

A suitable dosage for the vaccine for human infants is generally within the range of about 5 to 25 µg, or about 1 to 10 µg per kilogram of body weight.

5 EXAMPLES

The invention is illustrated by the following non-limiting examples. The N-acetyl, N-propionyl, N-butanoyl, N-isobutanoyl, N-pentanoyl and N-hexanoyl-GBMP-protein conjugates have been prepared for
10 evaluation purposes, and the results are discussed in the examples.

Materials and Methods for Preparing Conjugates

(a) Materials

15 Propionic, butanoic, isobutanoic, pentanoic, and hexanoic anhydrides together with colominic acid were obtained from Sigma Chemicals Co., St. Louis, MO. Because colominic acid is structurally identical to the group B meningococcal polysaccharide (GBMP), it is referred to henceforth as GBMP. Tetanus toxoid (TT) was obtained from the Institut Armand Frappier, Laval, Quebec, and its monomeric form, used in all the
20 conjugations, was obtained by passage of the above preparation through a Bio-Gel (trademark) A 0.5 (200-400 mesh) column (1.6 x 90 cm) (Bio-Rad, Richmond, CA.), equilibrated and eluted with 0.01 M phosphate buffered physiologic saline (PBS) (pH 7.4)

(b) N-Deacetylation of the GBMP

25 The GBMP (Na⁺ salt) (1.0g) was dissolved in 5 ml of 2M NaOH and, following the addition of NaBH₄ - (150 mg), the solution was heated at 110°C for 6 hours in a screw cap Teflon (trademark) container (60 mL). This procedure is essentially as described in J. Immunol., 134, 2651 (1985) and U.S. patent 4,727,136, both in the name of Harold J. Jennings, et al. The cooled diluted solution was then exhaustively dialyzed
30 against distilled water at 4°C, and lyophilized. The fact that N-deacetylated GBMP was obtained was determined by the absence of the methylacetamido signal (singlet at delta 2.07) in the ¹H-nmr spectrum of the N-deacetylated GBMP.

(c) N-Acylations of the GBMP

35 N-Deacetylated GBMP (1.0g) was dissolved in 50 mL of 5% aqueous NaHCO₃. To five individual aliquots (10 mL of the above solution) were added either propionic, butanoic, isobutanoic, pentanoic or hexanoic anhydrides. These reagents were added in 3 x 0.5 mL aliquots over a 3 hour period of time at room temperature while the solution was maintained at pH 8.0 with 0.5N NaOH. Methanol (0.5 mL) was
40 added simultaneously with each addition of anhydride in order to increase their solubility. Finally the solutions were stirred for 16 hours at 4°C, exhaustively dialysed against distilled water at 4°C, and lyophilized. The individual N-propionylated, N-butanoylated, N-isobutanoylated, N-pentanoylated and N-hexanoylated GBMP were all obtained in yields in excess of 90%. In each case, essentially complete N-acylation was confirmed by the disappearance in the respective ¹H-nmr spectrum of N-deacetylated GBMP.

(d) Sizing of the fragments of the different N-acylated GBMP

45 Gel filtration, using an Ultragel (trademark) AcA 44 (bead diameter 60-140 µm) column (IBF Biotechnics, Savage, MD.) with PBS as eluant, was employed to obtain the desired average molecular weight N-acylated GBMP material. Fractions eluting from the column at K_D 0.5 to K_D 0.7 as measured by FLPC (see
50 below) were collected, dialyzed, and lyophilized. This range of K_D values corresponds to N-acylated GBMP of approximately 30-50 sialic acid residues (10,000 to 15,000 Daltons, typically 12,000 Daltons average molecular weight). Fractions in the range of K_D 0.2 to 0.4 corresponding to fragments having an average molecular weight in the range of 30,000 to 40,000 Daltons have also been collected and conjugated. Thus,
55 N-acylated material eluting in the K_D range of 0.2 to 0.7 is of particular interest.

(e) Polysaccharide Conjugates

Terminal aldehyde groups were introduced into the N-acylated GBMP by periodate oxidation (see U.S. patent no. 4,356,170). The N-acylated GBMP fragments above were oxidized in 0.1 M aqueous NaIO₄ - (sodium metaperiodate) (10 mL) for 2 hours at room temperature in the dark. Excess periodate was then destroyed by the addition of 1 mL of ethylene glycol and the solution was then exhaustively dialyzed at 4°C, and lyophilized. The use of NaBH₄ in the N-deacetylation procedure (except for the GBMP) results in the transformation of the terminal reducing sialic acid residues of each of the N-acylated GBMP, to open chain polyol residues. This type of residue is periodate sensitive (see J. Immunol., 127, 1011 (1981) and U.S. patent 4,356,170 Harold J. Jennings et al), thereby resulting in the introduction of aldehyde groups into the N-acylated GBMP fragments at both termini.

The oxidized fragments (100 mg) were dissolved in 0.1 M NaHCO₃ (pH 8.1) buffer (2mL) and TT (20 mg) was added to the solution. Finally, following the addition of sodium cyanoborohydride (NaCNBH₃) (40 mg), the solution was gently stirred at room temperature. The course of the conjugation was followed by FPLC using a gel filtration column containing Superose (trademark) 12 HR10/30 (Pharmacia), run isocratically at 1 mL/min in PBS buffer at pH 7.2, both the protein and N-acylated GBMP fragments being monitored at 214 nm. The fragments had K_D 0.6, TT had K_D 0.39 and the conjugates had K_D 0.23. The conjugation was complete when all the TT was expended as determined by the loss of the peak in the FPLC chromatogram corresponding to the component at K_D 0.39. In most cases, the conjugations were complete in 2 days but were left for a total reaction time of 4 days. The potential unreacted aldehyde groups were finally reduced with sodium borohydride (20 mg) prior to gel filtration.

The polysaccharide - TT conjugates were separated from the polysaccharide fragments by gel filtration using a Bio Gel A column with PBS as eluant. The eluant containing the conjugate was dialyzed against distilled water and lyophilized. The N-acylated GBMP - TT conjugates contained from 12-30%, typically 12-20%, sialic acid as determined by the resorcinol method described by Svennerholm, L., Quantitative Estimation of Sialic Acids, II A Colorimetric Resorcinol-Hydrochloric Acid Method, Biochim. Biophys. Acta, 24, 604 (1957). This indicates that the conjugates had a molar ratio of polysaccharide to TT of 2-3:1 respectively.

30 Immunization and Immunoassays(a) Immunization Procedures

Twenty female white CFI mice (8-10 weeks old) were immunized intraperitoneally (3 times at 3 week intervals) with each individual N-acylated GBMP-TT conjugate in Freund's complete adjuvant (FCA) (Difco, Detroit, MI). Each immunization contained sufficient conjugate (10-12 µg) to contain 2 µg of sialic acid. Eleven days after the third injection, the mice were exsanguinated. The following tests were done on the sera.

(b) Radioactive antigen binding assay

This assay was carried out by a modification of the Farr technique using extrinsically [³H]-labeled GBMP (Jennings H.J., et al, Determinant Specificities of the Groups B and C polysaccharides of *Neisseria meningitidis*, J. Immunol., 134, 2651 (1985), or [³H]-labeled N-Pr-GBMP (Jennings H. J., et al, Unique Intermolecular Bactericidal Epitope involving the Homo-Sialo Polysaccharide Capsule on the Cell Surface of Group B *Neisseria meningitidis* and *Escherichia coli* K1, J. Immunol., 142, 3585-3591 (1989)). The reaction mixture for the radioactive antigen-binding assay was obtained by mixing in Eppendorf polypropylene micro test tubes 20 µL of pooled antisera, from groups of 20 mice immunized with each individual N-acylated GBMP-TT conjugate, diluted to 100 µL with PBS, with [³H]-labeled GBMP and [³H]-labeled N-Pr-GBMP in 50 µL of PBS. After incubation at 4°C for 16 hours, 150 µL of saturated (at 4°C) ammonium sulfate (pH 7.0) was added to the tubes and the tubes agitated and left to stand at 4°C for 30 min. The tubes were centrifuged at 15,000 rpm for 10 min. and two aliquots of 130 µL were drawn from the tubes. The aliquots were mixed with 2 mL of water and a scintillant-containing xylene (ACS aqueous scintillant) and the mixtures were counted in a liquid scintillation counter. Results are given in Table 1.

Table 1

5	Binding of [³ H] - labeled - N-Ac-GBMP to different mouse anti- <u>N</u> - acyl - GBMP-TT conjugate sera.				
	Antiserum	% Binding ^a			
		1	2	3	4
	<u>N</u> -Pr-GBMP-TT	41	40	39	12
	<u>N</u> -Bu-GBMP-TT	4	4	7	4
10	<u>N</u> -IsoBu-GBMP-TT	9	-	-	-
	<u>N</u> -Pen-GBMP-TT	36	-	-	-
	<u>N</u> -Hex-GBMP-TT	16	-	-	-

^a The four binding experiments were carried out on pooled antisera from 20 immunized mice.

Abbreviations used in Table 1 and other tables:

15 N-Ac-, N-Pr, N-Bu N-IsoBu, N-Pen, N-Hex stand for N-Acetyl, N-Propionyl-, N-Butanoyl-, N-Isobutanoyl-, N-Pentanoyl- and N-Hexanoyl-.

20 The numerals 1, 2, 3 and 4 are results of four repeat experiments. Table 1 demonstrates conclusively that the N-Ac-GBMP (which carries the same epitope as fetal N-CAM) binds less to the antiserum induced by the N-Bu-GBMP, N-IsoBu-GBMP, N-Pen-GBMP and N-Hex-GBMP than that induced by the N-Pr-GBMP. From this, it can be deduced from Table 1 that the N-Bu-, N-IsoBu, N-Pen- and N-Hex-polysaccharide-conjugates raise less cross-reactive antibodies than the N-Pr-conjugate.

(c) Quantitative precipitin analyses

25 These experiments were carried out by the method of Kabat and Mayer, Experimental Immunochimistry Charles C. Thomas, Springfield, p.22 (1961). Aliquots (100 μ L) of anti-N-acyl GBMP-TT sera (diluted 5 fold in PBS) were reacted in tubes with increasing concentrations of the N-acetyl (colominic acid), N-propionyl, N-butanoyl, N-isobutanoyl, N-pentanoyl and N-hexanoyl GBMP in a total volume of 200 μ L (adjusted with PBS). The higher molecular weight fractions of these derivatives were used in these experiments and they were obtained from the eluate of the Ultragel Aca 44 column (K_D 0.4 as measured by FPLC) previously used to size the fragments of the N-acylated GBMP. The tubes were incubated at 4° C for 4 days with daily mixing, centrifuged, and the quantity of antibody protein was determined by the method of Lowry et al, Protein Measurement with the Folin phenol reagent, J. Biol. Chem., 1933, 265 (1951). The results are given in Table 2.

Table 2

40	Precipitation ^a of mouse anti- <u>N</u> -acyl-GBMP-TT sera using different <u>N</u> -acyl GBMP as precipitating antigens.					
	Antiserum	<u>N</u> -acyl-GBMP antigen				
		<u>N</u> -acetyl	<u>N</u> -propionyl	<u>N</u> -butanoyl	<u>N</u> -pentanoyl	<u>N</u> -hexanoyl
	<u>N</u> -Pr-GBMP-TT	0.16	0.40	0.20	0.15	0.15
	<u>N</u> -Bu-GBMP-TT	0.04	1.15	2.60	3.20	1.90
45	<u>N</u> -Pen-GBMP-TT	0.13	0.38	0.44	6.35	3.55
	<u>N</u> -Hex-GBMP-TT	0.02	0.08	0.80	4.15	4.40

^a Maximum amount of antibody precipitated expressed in mg/mL of antiserum

50 As regards cross-reactivity, the first column of Table 2 indicates that very little cross-reactive antibodies are produced by the N-Bu and N-Hex conjugates as compared to the N-Pr conjugate. It can also be seen that the N-Pen conjugate produces less cross-reactive antibodies than the N-Pr conjugate.

With reference to immunogenicity, in terms of homologous response, it can be seen from Table 2 that the N-Bu-(2.60), N-Pen- (6.35) and N-Hex- (4.40) GBMP-TT conjugates are more immunogenic than the N-Pr-GBMP analog (0.40).

(d) ELISA

55 The wells of EIA microtitration plates (Flow Labs, Mississauga, Ontario, Canada) were coated with a 10 μ g/mL solution of either GBMP-, NPrGBMP- or NBu-GBMP-BSA conjugates in PBS (100 μ L/well).

The plates were left for 18 hours at 4°C and after coating they were washed with 1% bovine serum albumin in PBS for 10 min. at room temperature (blocking step). The wells were then filled with 100 µL of serial 10-fold dilutions in PBS of anti-mouse-N-acyl GBMP-TT conjugate sera and the plates were incubated for 1 hour at room temperature. After washing with SBT the plates were incubated for 1 hour at room temperature with 50 µL of the appropriate dilution of goat antimouse immunoglobulin peroxidase labeled conjugates (Kirkegard and Perry Laboratories), then the contents of the wells were aspirated and the plates washed five times with SBT. Finally 50 µL of Tetramethylene blue-peroxidase substrate (TMB) (Kirkegard and Perry Laboratories) were added to each well and after 10 min the absorbance at 450 nm was measured with a Multiscan spectrophotometer (Flow Laboratories, Mississauga, Ont.) Results are given in Table 3.

Table 3

ELISA titrations of pooled mouse anti-N-acyl-GBMP-TT conjugate serum against N-acyl-GBMP-BSA conjugates.			
Coating Antigen	Titers ^a of Antisera		
	a. N-Pr-GBMP ^b	a. N-Bu-GBMP ^b	a. N-IsoBu-GBMP ^b
N-Ac-GBMP-BSA	7800	1000	7000
N-Pr-GBMP-BSA	40000	39000	9800
N-Bu-GBMP-BSA	26000	52000	9700
N-IsoBu-GBMP-BSA	-	-	25000

a titer (GM) = reciprocal of dilution at 50% of the maximum absorbance at 450 nm.

b N-acyl specific antisera induced in mice by homologous N-acyl-GBMP-TT conjugates.

With reference to cross-reactivity, it can be seen from Table 3 that the N-Bu-GBMP-TT conjugate raises less cross-reactive antibodies with respect to N-Ac-GBMP (1000) than does the N-Pr-GBMP-TT conjugate (7800). The reason for this is that the GBMP binds less to antibody induced by the N-Pr-GBMP-TT conjugate than that induced by the N-Bu-GBMP-TT conjugate. Similar comments apply with respect to the N-IsoBut-GBMP-TT conjugate.

As regards immunogenicity, the N-Bu conjugate is more immunogenic than the N-Pr analogue, as shown by the homologous binding titers of 52,000 (N-Bu) and 40,000 (N-Pr).

(e) Radioactive binding inhibition assay Increasing concentration of the larger molecular sized N-acyl GBMP inhibitor in PBS (80 µL) were added to 20 µL of mouse anti-N-Pr-GBMP-TT conjugate antiserum, an amount sufficient to bind 50% of the (³H)-labeled N-Pr-GBMP in the absence of inhibitor. The tubes were incubated for 1 hour at 37°C and 50 µL of (³H)-labeled N-Pr-GBMP in PBS was added. After gentle mixing the tubes were incubated at 4°C for 16 hours and the assays were performed exactly as described previously for the radioactive antigen binding assay. The % inhibition was calculated using the following formula:

$$\text{percent inhibition} = 100 \times [(\text{cpm with inhibitor minus cpm without inhibitor}) / (\text{cpm without antibody minus cpm without inhibitor})]$$

Results are given in Table 4.

Table 4

Inhibition of binding of [³ H] - labeled N-Pr-GBMP to mouse anti-N-Pr-GBMP-TT conjugate induced IgG _{2a} , IgG _{2b} (A) ^a and IgG ₁ (B) ^a antibodies.		
Inhibitor ^b	A	B
N-Ac-GBMP	>50.0	>50.0
N-Pr-GBMP	0.6	0.3
N-Bu-GBMP	0.3	0.2
N-IsoBu-GBMP	>50.0	-
N-Pen-GBMP	2.3	2.5
N-Hex-GBMP	10.2	10.0

a These were fractions of mouse polyclonal anti-N-Pr-GBMP-TT, serum, described in Jennings et al, J. Immunol., 142, 3585-3591 (1989).

b Micrograms of inhibitor to give 50% inhibition.

Bactericidal assays These assays were carried out by the method described by Jennings et al, J. Exp. Med., 165, 1207-1211 (1987).

Neisseria meningitidis strain B (M 986) was used in these assays. Results are given in Table 5.

Table 5

Binding of [³ H] - labeled - N-Pr - GBMP to different mouse anti - N-acyl - GBMP-TT conjugate sera and the bactericidal titers of the respective antisera.		
Antiserum	μL of antiserum ^a	Bactericidal Titer ^b
N-Pr-GBMP-TT	13	128
N-Bu-GBMP-TT	10	64
N-IsoBu-GBMP-TT	ND	64
N-Pen-GBMP-TT	24	64
N-Hex-GBMP-TT	>100	8
N-Ac-GBMP-TT	>100	8

a μL of antiserum (diluted 5 fold with PBS) required for 50% binding.

b Dilution experiment: one dilution difference, e.g. 128 as compared to 64, is within experimental error.

Table 4 illustrates that N-Bu-GBMP is as good an inhibitor as the N-Pr-GBMP for the binding of the latter to its own homologous antibodies. Therefore, the use of N-Bu-GBMP in place of N-Pr-GBMP does not result in any significant loss of properties exhibited by N-Pr-GBMP. Table 5 demonstrates that the N-Bu-GBMP binds as well as the N-Pr-GBMP to antibodies induced by the N-Pr-GBMP-TT conjugate. Antisera induced by both the N-Bu-GBMP-TT and N-Pr-GBMP-TT conjugates gave similar bactericidal titers. This evidence indicates the equivalence of the N-Bu-, N-IsoBu- and N-Pen-GBMP to the N-Pr-GBMP in their ability to mimic the bactericidal epitope on the surface of the group B meningococci.

Claims

1. A modified group B polysaccharide of *Neisseria meningitidis* wherein the sialic acid residue N-acetyl groups of the native polysaccharide are replaced by C₄-C₈-acyl groups, wherein the modified polysaccharide has an average molecular weight within the range of 10,000 to 50,000 Daltons.
2. A modified *E. coli* K1 capsular polysaccharide wherein the sialic acid residue N-acetyl groups of the native polysaccharide are replaced by a C₄-C₈-acyl group, wherein the modified polysaccharide has an average molecular weight within the range of 10,000 to 50,000 Daltons.

3. A modified polysaccharide according to claim 1, wherein the C₄-C₈-acyl group is selected from the group consisting of n-butanoyl, iso-butanoyl, n-pentanoyl, iso-pentanoyl, neo-pentanoyl, hexanoyl, heptanoyl and octanoyl.
- 5 4. A modified polysaccharide according to claim 3, wherein the acyl group is selected from the group consisting of n-butanoyl, iso-butanoyl, n-pentanoyl and hexanoyl.
5. A modified polysaccharide according to claim 2, wherein the C₄-C₈-acyl group is selected from the group consisting of n-butanoyl, iso-butanoyl, n-pentanoyl, iso-pentanoyl, neo-pentanoyl, hexanoyl, heptanoyl and octanoyl.
- 10 6. A modified polysaccharide according to claim 5, wherein the acyl group is selected from the group consisting of n-butanoyl, iso-butanoyl, n-pentanoyl and hexanoyl.
- 15 7. The polysaccharide according to claim 1 wherein about 27% to 100% of the sialic acid residue N-acetyl groups have been replaced by C₄-C₈-acyl groups.
8. The polysaccharide according to claim 2 wherein about 27% to 100% of the sialic acid residue N-acetyl groups have been replaced by C₄-C₈-acyl groups.
- 20 9. An antigenic conjugate comprising a modified group B polysaccharide of *Neisseria meningitidis* having sialic acid residue N-acetyl group replaced by a N-C₄-C₈-acyl group conjugated to an immunologically-suitable protein, wherein the modified polysaccharide has an average molecular weight within the range of 10,000 to 50,000 Daltons.
- 25 10. A conjugate according to claim 9, wherein the acyl group is selected from the group consisting of n-butanoyl, iso-butanoyl, n-pentanoyl, iso-pentanoyl, neo-pentanoyl, hexanoyl, heptanoyl and octanoyl.
11. A conjugate according to claim 10, wherein the acyl group is selected from the group consisting of n-butanoyl, iso-butanoyl, n-pentanoyl and hexanoyl.
- 30 12. A conjugate according to claim 9, wherein the protein is selected from the group consisting of tetanus toxoid, diphtheria toxoid, a cross-reacting material (CRM) and a bacterial protein carrier.
13. A conjugate according to claim 9, wherein the protein and the polysaccharide are covalently linked through a -CH₂-NH-protein linkage.
- 35 14. The conjugate according to claim 12, wherein the CRM is CRM₁₉₇.
- 40 15. A vaccine comprising a conjugate of claim 9 and an pharmaceutically acceptable injectable carrier.
16. A vaccine according to claim 15, and further comprising an physiologically acceptable adjuvant.
17. A vaccine according to claim 16, wherein said adjuvant is selected from the group consisting of aluminum hydroxide, aluminum phosphate and aluminum sulphate.
- 45 18. A gamma globulin fraction capable of passive protection against meningitis caused by group B *N. meningitidis* and *E. coli* K1, said gamma globulin fraction being produced by immunizing a mammal with a vaccine as claimed in Claim 17, and recovering gamma globulin from said mammal.
- 50 19. The modified polysaccharide according to claim 1 wherein about 90% to 100% of the sialic acid residue N-acetyl groups are replaced by C₄ - C₈ acyl groups.
20. The polysaccharide according to claim 2 wherein about 90% to about 100% of the sialic acid residue N-acetyl groups are replaced by C₄ - C₈ acyl groups.
- 55

Patentansprüche

1. Modifiziertes Polysaccharid der Gruppe B von *Neisseria meningitidis*, worin die N-Acetylgruppen des Sialinsäurerestes des nativen Polysaccharids durch C₄-C₈-Acylgruppen ersetzt sind, wobei das modifizierte Polysaccharid ein durchschnittliches Molekulargewicht im Bereich von 10 000 bis 50 000 Dalton hat.
2. Modifiziertes *E.coli*-K1-Kapselpolysaccharid, worin die N-Acetylgruppen des Sialinsäurerestes des nativen Polysaccharids durch eine C₄-C₈-Acylgruppe ersetzt sind, wobei das modifizierte Polysaccharid ein durchschnittliches Molekulargewicht im Bereich von 10 000 bis 50 000 Dalton hat.
3. Modifiziertes Polysaccharid nach Anspruch 1, wobei die C₄-C₈-Acylgruppe aus der aus n-Butanoyl, iso-Butanoyl, n-Pentanoyl, iso-Pentanoyl, neo-Pentanoyl, Hexanoyl, Heptanoyl und Octanoyl bestehenden Gruppe gewählt ist.
4. Modifiziertes Polysaccharid nach Anspruch 3, wobei die Acylgruppe aus der aus n-Butanoyl, iso-Butanoyl, n-Pentanoyl und Hexanoyl bestehenden Gruppe gewählt ist.
5. Modifiziertes Polysaccharid nach Anspruch 2, wobei die C₄-C₈-Acylgruppe aus der aus n-Butanoyl, iso-Butanoyl, n-Pentanoyl, iso-Pentanoyl, neo-Pentanoyl, Hexanoyl, Heptanoyl und Octanoyl bestehenden Gruppe gewählt ist.
6. Modifiziertes Polysaccharid nach Anspruch 5, wobei die aus der aus n-Butanoyl, iso-Butanoyl, n-Pentanoyl und Hexanoyl bestehenden Gruppe gewählt ist.
7. Polysaccharid nach Anspruch 1, worin etwa 27 % bis 100 % der N-Acetylgruppen des Sialinsäurerestes durch C₄-C₈-Acylgruppe ersetzt sind.
8. Polysaccharid nach Anspruch 2, worin etwa 27 % bis 100 % der N-Acetylgruppen des Sialinsäurerestes durch C₄-C₈-Acylgruppe ersetzt sind.
9. Antigenes Konjugat, umfassend ein modifiziertes Polysaccharid der Gruppe B von *Neisseria meningitidis*, bei dem die N-Acetylgruppen des Sialinsäurerestes durch eine N-C₄-C₈-Acylgruppe ersetzt ist, konjugiert mit einem immunologisch geeignetem Protein, wobei das modifizierte Polysaccharid ein durchschnittliches Molekulargewicht im Bereich von 10 000 bis 50 000 Dalton besitzt.
10. Konjugat nach Anspruch 9, wobei die Acylgruppe aus der aus n-Butanoyl, iso-Butanoyl, n-Pentanoyl, iso-Pentanoyl, neo-Pentanoyl, Hexanoyl, Heptanoyl und Octanoyl bestehenden Gruppe gewählt ist.
11. Konjugat nach Anspruch 10, wobei die Acylgruppe aus der aus n-Butanoyl, iso-Butanoyl, n-Pentanoyl und Hexanoyl bestehenden Gruppe gewählt ist.
12. Konjugat nach Anspruch 9, wobei das Protein aus der aus Tetanus-Toxoid, Diphtherie-Toxoid, einem kreuzreagierenden Material (CRM) und einem bakteriellen Proteinträger bestehenden Gruppe gewählt ist.
13. Konjugat nach Anspruch 9, wobei das Protein und das Polysaccharid durch eine -CH₂-NH-Proteinbindung kovalent verknüpft sind.
14. Konjugat nach Anspruch 12, bei dem das CRM CRM₁₉₇ ist.
15. Impfstoff, der ein Konjugat nach Anspruch 9 und einen pharmazeutisch annehmbaren injizierbaren Träger umfaßt.
16. Impfstoff nach Anspruch 15, der ferner ein physiologisch annehmbares Adjuvans umfaßt.
17. Impfstoff nach Anspruch 16, bei dem das Adjuvans aus der aus Aluminiumhydroxid, Aluminiumphosphat und Aluminiumsulfat bestehenden Gruppe gewählt wird.

18. Gamma-Globulinfraktion, die zum passiven Schutz gegen durch *N. meningitidis* der Gruppe B und *E.coli* K1 verursachte Meningitis in der Lage ist, wobei die gamma-Globulinfraktion hergestellt wird, indem ein Säugetier mit einem Impfstoff nach Anspruch 17 immunisiert und das gamma-Globulin aus dem Säugetier gewonnen wird.
19. Modifiziertes Polysaccharid nach Anspruch 1, worin etwa 90 % bis 100 % der N-Acetylgruppen des Sialinsäurerestes durch C₄-C₈-Acylgruppe ersetzt sind.
20. Polysaccharid nach Anspruch 2, worin etwa 90 % bis etwa 100 % der N-Acetylgruppen des Sialinsäurerestes durch C₄-C₈-Acylgruppe ersetzt sind.

Revendications

1. Polysaccharide de *Neisseria meningitidis* du groupe B, modifié, dans lequel les groupements N-acétyles des résidus d'acide sialique du polysaccharide natif sont remplacés par des groupements acyles en C₄-C₈, dans lequel le polysaccharide modifié possède une masse moléculaire moyenne comprise dans la gamme allant de 10 000 à 50 000 Daltons.
2. Polysaccharide capsulaire de *E. coli* K1, modifié, dans lequel les groupements N-acétyles des résidus d'acide sialique du polysaccharide natif sont remplacés par un groupement acyle en C₄-C₈, dans lequel le polysaccharide modifié possède une masse moléculaire moyenne comprise dans la gamme allant de 10 000 à 50 000 Daltons.
3. Polysaccharide modifié selon la revendication 1, dans lequel le groupement acyle en C₄-C₈ est choisi dans le groupe constitué par le n-butanoyle, l'isobutanoyle, le n-pentanoyle, l'isopentanoyle, le néopentanoyle, l'hexanoyle, l'heptanoyle et l'octanoyle.
4. Polysaccharide modifié selon la revendication 3, dans lequel le groupement acyle est choisi dans le groupe constitué par le n-butanoyle, l'isobutanoyle, le n-pentanoyle, et l'hexanoyle.
5. Polysaccharide modifié selon la revendication 2, dans lequel le groupement acyle en C₄-C₈ est choisi dans le groupe constitué par le n-butanoyle, l'isobutanoyle, le n-pentanoyle, l'isopentanoyle, le néopentanoyle, l'hexanoyle, l'heptanoyle et l'octanoyle.
6. Polysaccharide modifié selon la revendication 5, dans lequel le groupement acyle est choisi dans le groupe constitué par le n-butanoyle, l'isobutanoyle, le n-pentanoyle, et l'hexanoyle.
7. Polysaccharide selon la revendication 1, dans lequel environ 27% à 100% des groupements N-acétyles des résidus d'acide sialique ont été remplacés par des groupements acyles en C₄-C₈.
8. Polysaccharide selon la revendication 2, dans lequel environ 27% à 100% des groupements N-acétyles des résidus d'acide sialique ont été remplacés par des groupements acyles en C₄-C₈.
9. Conjugué antigénique comprenant un polysaccharide de *Neisseria meningitidis* du groupe B, modifié, ayant un groupement N-acétyle de résidu d'acide sialique, remplacé par un groupement N-acyle en C₄-C₈, conjugué à une protéine immunologiquement convenable, dans lequel le polysaccharide modifié possède une masse moléculaire moyenne comprise dans la gamme allant de 10 000 à 50 000 Daltons.
10. Conjugué selon la revendication 9, dans lequel le groupement acyle est choisi dans le groupe constitué par le n-butanoyle, l'isobutanoyle, le n-pentanoyle, l'isopentanoyle, le néopentanoyle, l'hexanoyle, l'heptanoyle et l'octanoyle.
11. Conjugué selon la revendication 10, dans lequel le groupement acyle est choisi dans le groupe constitué par le n-butanoyle, l'isobutanoyle, le n-pentanoyle, et l'hexanoyle.
12. Conjugué selon la revendication 9, dans lequel la protéine est choisie dans le groupe constitué par la toxoïde tétanique, la toxoïde diphtérique, une matière à réaction croisée (CRM) et un porteur protéique d'origine bactérienne.

13. Conjugué selon la revendication 9, dans lequel la protéine et le polysaccharide sont liés de manière covalente par l'intermédiaire d'une liaison $-CH_2-NH$ -protéine.

14. Conjugué selon la revendication 12, dans lequel la CRM est la CRM₁₉₇.

5

15. Vaccin comprenant un conjugué selon la revendication 9 et un véhicule injectable pharmaceutiquement acceptable.

16. Vaccin selon la revendication 15, et comprenant en outre un adjuvant physiologiquement acceptable.

10

17. Vaccin selon la revendication 16, dans lequel ledit adjuvant est choisi dans le groupe constitué par l'hydroxyde d'aluminium, le phosphate d'aluminium et le sulfate d'aluminium.

18. Fraction de gammaglobuline procurant une protection passive contre la méningite causée par le N. meningitidis du groupe a et E. coli K1, ladite fraction gammaglobuline étant produite en immunisant un mammifère avec un vaccin tel que revendiqué à la revendication 17, et en récupérant la gammaglobuline dudit mammifère.

15

19. Polysaccharide modifié selon la revendication 1, dans lequel environ 90% à 100% des groupements N-acétyles des résidus d'acide sialique sont remplacés par des groupements acyles en C₄-C₈.

20

20. Polysaccharide selon la revendication 2, dans lequel environ 90% à environ 100% des groupements N-acétyles des résidus d'acide sialique sont remplacés par des groupements acyles en C₄-C₈.

25

30

35

40

45

50

55